

Occurrence of intestinal and urinary parasites among Alkhalawi students, sudan: a cross-sectional study

Comment [u1]: Intestinal and urine parasite prevalence in Alkhalawi students, Sudan: a cross-sectional investigation

abstract

Background: Parasitic diseases were some of the leading causes of global mortality with higher burdens of prevalence in developing countries, especially regions of the world where the contaminated water and poor personal hygiene. The purpose of this study to determine the prevalence rate of intestinal and urinary parasites among Alkhalawi students in Rabak city, White Nile State, Sudan.

Materials and methods: In this study, 150 stool and 150 urine samples were collected from Alkhalawi students (all were males). The stool samples were processed by wet preparation, formal ether concentration technique and the urine samples were processed by the sedimentation technique.

results: The results revealed that out of the 150 stool and 150 urine samples, 37 were found positive for intestinal and urinary parasites, this constituted an overall prevalence rate of 12.3%. The highest prevalence rate (28%) was reported among the 11-15 years age groups, while the lowest prevalence rate (0%) was reported among the 21-25 years age groups. The prevalence rate of different parasites was as follow: *Giardia lamblia* (11.3%), *Entamoebahistolytica* (8%), *Hymenolepis nana* (2.7%), *Schistosomahaematobium* (3.3%). The prevalence of intestinal parasites (21.3%) was higher than urinary parasites (3.3%). The study showed that 24 % were infected with single infection and 0.6% were infected with mixed infection.

conclusion: From the result we concluded the intestinal and urinary parasites are highly prevalent among the Alkhalawi students in Rabak city. *Giardia lamblia* is more likely to occur among the Alkhalawi students followed by *Entamoebahistolytica*. The highest infection rate was reported among the 11-15 years age groups. Formal ether concentration technique proved to be as efficient more than the wet preparation technique.

Keywords: intestinal parasites; urinary parasites; prevalence rate; Alkhalawi students; Sudan

Comment [u2]: The abstract effectively communicates the purpose, methodology, and key findings of the study on the prevalence of intestinal and urinary parasites among Alkhalawi students in Rabak city, Sudan.

Comment [u3]: Reduce overall abstract to 250 words from to 280 pls

Comment [u4]: Provide pls

Keywords

introduction

Infection with intestinal parasites remains one of the most common health problems in the world mainly in tropical and subtropical countries [1]. Parasitic infection associated with poor hygiene and lack of sanitation is commonly observed in developing countries [2]. Intestinal and urinary parasites are distributed worldwide, particularly in the tropical and subtropical areas [3]. The prevalence of these parasites is promoted by several epidemiological factors such as poor sanitation, environmental degradation, ignorance, poor personal and community hygiene, climate condition and other socio-cultural practice such as the use of night soil as fertilizer [4]. The intestinal parasitic infections (e.g.

helminths and pathogenic intestinal protozoa) are of public health importance, particularly in developing countries. For example, the global burden caused by soil transmitted helminthiasis is estimated at 39 million disability-adjusted life years (DALYs) [5]. Whereas urinary schistosomiasis being a major debilitated disease characterized by blood in the urine. In the worst cases, schistosomiasis will cause bladder cancer, caused by the parasite *Schistosomahaematobium*, about 38 million people are infected in 16 African countries [6]. Intestinal parasites spread rapidly in areas with poor sanitation and are the most common in tropical developing countries on the African, Asia and South American continents [7]. Intestinal parasites and

Comment [u5]: Write in Upper Case

urinary parasite infection are a common health problem especially among children and communities with poor hygiene and unsafe water supply. The infections with intestinal and urinary parasites are serious and may lead to death. The infections are directly related to the behavior of the infected person. Therefore, the purpose of this study to determine the prevalence rate of intestinal and urinary parasites among Alkhalawi students in Rabak city, White Nile State, Sudan.

Materials and methods

This was a cross-sectional study. The study was performed in four Islamic centers (Alkhalawi) in Rabak in the White Nile State. The study was conducted on male students in the Islamic centers (Alkhalawi) in Rabak city. The population was divided according to age groups as follow: Group A from 5 – 10, Group B from 11 – 15, Group C from 16 – 20, and Group D from 21– 25 years-old.

ethics approval and consent to participate

Ethical approval of the study was given from ALNeelain University, college of graduate studies and the informed consents were taken from the all study subjects after explaining the nature of the study.

samples reception and processing

Sterile plastic containers were distributed which were labeled with the name and an assigned code to collect the samples; information and explanations were furnished to how they were to collect and transport the stool and urine samples to the collection sites. The collection and transportation guidelines followed pre-analytical standard operating procedures (SOPs). 150 stool and 150 urine samples were collected in this study according to the formula $(n=z^2pq/d)$.

Detection of intestinal parasites

Wet preparation and formal-ether concentration technique were used to detect and confirm the intestinal parasites.

Detection of urinary parasites

Urine sedimentation technique was used to detect the urinary parasites.

Inclusion criteria: The Alkhalawi students in Rabak city.

Exclusion criteria: Population other than the Alkhalawi students.

statistic

Data collected were analyzed by using statistical package for the social sciences (SPSS) software. Statistical analysis was done using Chi-square to evaluate any association between parasitic infections. Observed differences in data were considered significant and noted in the text if $P < 0.05$. Data were presented in tables.

results

Overall prevalence rate of intestinal and urinary parasites among the Alkhalawi students in Rabak city, White Nile State

The results showed that out of the 150 stool and 150 urine samples of the Alkhalawi students examined, 37 were found positive for gastrointestinal and urinary parasites. This constituted an overall prevalence rate of 12.3% (Table 1).

Table 1: Overall prevalence rate of intestinal and urinary parasites among the Alkhalawi students in Rabak city, White Nile State.

Number of examined subjects	Number of positive subjects	Prevalence %
300	37	12.3

Comparison between intestinal parasites infection among the study population

The study revealed that out of the 32 positive intestinal parasites cases, 31 were with a single infection and only one was with a mixed infection (*E.histolytica* and *G.lambliia*) constituting a prevalence rates of 20.6% and 0.6% respectively (Table 2). The difference in rates was found to be statistically significant at $p=0.000$.

Comparison between the prevalence rates of intestinal parasites using direct wet preparation and formal ether concentration technique

Comment [u12]: Statistical Analysis

Comment [u6]: State clearly, aim of the research, objectives and hypotheses

Comment [u7]: Write in upper case

Comment [u13]: The statistical package for the social sciences (SPSS) was used to evaluate the data that were gathered. Chi-square statistical analysis was used to assess any potential correlations between parasite illnesses. If $P < 0.05$, then the observed differences in the data were deemed significant and reported in the text.

Comment [u14]: Write in Upper case

Comment [u8]: Recast this way: This investigation was cross-sectional. Four Islamic centers in Rabak, the White Nile State (Alkhalawi), served as the study's locations. Male students in Rabak City's Islamic centers (Alkhalawi) were the subjects of the study. The age groupings of the population were as follows: Group A consisted of people aged 5 to 10, Group B of people aged 11 to 15, Group C of people aged 16 to 20, and Group D of people aged 21 to 25.

Comment [u9]: Sample collection

Comment [u10]: Provide reference?

Comment [u11]: Intestinal parasite identification

UNDER PEER REVIEW

The prevalence of intestinal parasites by different parasitological techniques was as follows: 23% by direct wet preparation and 24.6% by formal ether (Table 2). The difference in rates was found to be statistically insignificant at $p=0.892$.

Comment [u18]: Why is that so, state your reasons then back with statistic p-value

Table 2: Comparison between the prevalence rates of intestinal parasites according to the infection type and used technique.

Comment [u15]: All table titles in title case

Character		Examined number	Positive samples	Percent %
Infection	Single infection	150	31	20.6
	Mixed infection (E.histolytica + G.lamblia)	150	1	0.6
Technique	Direct wet preparation	150	35	23
	Concentration technique	150	37	24.6

The prevalence rate of urinary and intestinal parasites among the Alkhalawi students in Rabak city, White Nile State according to age groups

Different intestinal and urinary parasites detected among the Alkhalawi students in Rabak city, White Nile State

The results showed that 5-10, 11-15, 16-20 and 21-25-years age group revealed different rates (13%, 28%, 18% and 0% respectively) (Table 3). The difference in rates was found to be statistically

The prevalence of different parasites was as follows: *Giardia lamblia* (11.3%), *Entamoebahistolytica* (8%), *Hymenolepis nana* (2.7%) and *Schistosoma haematobium* (3.3%) (Table 3).

Comment [u16]: According to Table 3, the findings indicated that the age groups of 5-10, 11-15, 16-20, and 21-25 years old showed varying rates (13%, 28%, 18%, and 0%, respectively). It was discovered that the rate differential was significantly

insignificant at $p=0.207$.

Table 3: The prevalence rate of urinary and intestinal parasites among the Alkhalawi students in Rabak city, White Nile State according to the age groups and detected parasite.

Comment [u17]: Write in title case

	Character	Examined Number	Positive Samples	Percent %
Age groups	A (5-10)	23	3	13
	B (11-15)	104	30	28
	C (16-20)	22	4	18
	D (21-25)	1	0	0
The parasite	<i>Giardia lamblia</i>	150	17	11.3
	<i>Entamoebahistolytica</i>	150	12	8
	<i>Hymenolepis nana</i>	150	4	2.7
	<i>Schistosoma haematobium</i>	150	5	3.3

UNDER PEER REVIEW

Discussion

Unsafe water and poor sanitation and hygiene have been reported to rank third among the 20 leading risk factors for health burden in developing countries, including Sudan [8]. From the results, it was obvious that the overall prevalence of intestinal and urinary parasites among the Alkhalawi students in Rabak city, White Nile state was high (12.3%). This rate was found to be lower than the rate reported by Akinboye *et al* (2015)[9] in Nigeria (53.3%). In this study, the highest prevalence rate (28%) was reported among the 11-15 years age group. This rate does not agree with Abuobieda (2017)[10] who reported that the age group 6-8 years was the most affected. The finding of this study indicated that the common intestinal and urinary parasites among the Alkhalawi students were: *Giardia lamblia* (11.3%), *Entamoebahistololytica* (8%), *Hymenolepis nana* (2.7%), *Schistosomahaematobium* (3.3%), while Colman *et al* (2013)[11] in Maiduguri reported that the most prevalent intestinal and urinary parasites were *Entamoeba coli* (9.95%), Hookworms (6.47%), *Entamoebahistololytica* (4.48%), *Hymenolepis nana* (3.48%), *Schistosomahaematobium* (3.48%) and *Taeniaspp* (0.05%). As far as the detection rates for the two techniques used, it was obvious that the detection rate (24.6%) was reported for the formal ether concentration technique and 23% for the wet preparation, which indicates that both techniques have the same efficacy in detecting intestinal parasites. Our results, for the formal ether technique does not agree with Venkatesh (2016) [12], in Hyderabad who reported 66% detection rate.

conclusion

From the result we can conclude that the intestinal and urinary parasites are highly prevalent among the Alkhalawi students in Rabak city. *Giardia lamblia* is more likely to occur among the Alkhalawi students in Rabak city followed by *Entamoebahistololytica*. The highest infection rate was reported among the 11-15 years age groups. Formal ether concentration technique proved to be as efficient more than the wet preparation technique. The Alkhalawi students are targeted to be informed and equipped with the needed information to avoid these infections.

References

- [1] Wakid MH. Improvement of Ritchie technique by identifying the food that can be consumed pre- analysis. J ApplSci Res. 2009; 5(3):293–296.
- [2] Wegayehu TT, Tsalla B, Seifu B, Teklu T. Prevalence of intestinal parasitic infections among highland and lowland dwellers in Gamo area, South Ethiopia. BMC Public Health. 2013; 13:151.
- [3] Ochei OJ, Kolhatkar A. Laboratory techniques for intestinal parasites. Medical laboratory science theory and practical. 2000 ; 909–1044.
- [4] Eneanya CI, Njom VS. Geohelminth contamination of some common fruits and vegetable in Enugu, South east Nigeria. The Niger J Parasitol. 2003; 24(1):123–128.
- [5] WHO. Prevention and control of schistosomiasis and soil transmission helminthiasis: first report of the joint WHO expert committees. WHO Tech Rep Ser. 2002; 912:1–57.
- [6] WHO. Schistosomiasis. World Health Organization division of control of tropical diseases January report series. 2007; pp.86.
- [7] Haftu D, Deyessa N, Agedew E. Prevalence and determinant factors of intestinal factors of intestinal parasites among school children in Arba Minch Town, Southern Ethiopia. Am J Health Res. 2014; 2(5):247–254.
- [8] World Health Organization. Emerging issues in water and infectious disease, 2003.
- [9] Akinboye DO, Abdullah AR, Awodele O, Akintunde TI, Effedua HI, et al. Prevalence of intestinal and urinary parasites among food-handlers in Ilishan-Remo, Ogun State, Nigeria. Nigerian Journal of Parasitology 2015; 36(2):113–118.
- [10] Abuobieda SMS. The prevalence rate of intestinal parasites among primary schools' children in Ombda locality, 2017.
- [11] Colman S, Mangoro ZM, Isa L. Incidence of intestinal and urinary parasites among prison inmates. AJMR. 2013; 1(1):11–15.
- [12] Venkatesh BS, Rao SR, Vivekan N, Shanker BM. A comparative study of concentration techniques for detection of intestinal parasitic infections—to evaluate the prevalence and to identify a better method of concentration technique at a Tribal Tertiary Care Hospital, 2016.

Comment [u19]: 11-15 years

Comment [u20]: Based on the findings, we may infer that Alkhalawi students in Rabak City have a significant prevalence of intestinal and urinary parasites. In Rabak, students who identify as Alkhalawi are more likely to contract *Giardia lamblia*, with *Entamoebahistololytica* following closely after. The age range of 11-15 years had the highest infection rate recorded. The wet preparation method was found to be less effective than the formal ether concentration method. The goal is to provide the Alkhalawi students with the knowledge and tools they need to prevent contracting these infections.