

VESICULAR NANOCARRIERS IN TRANSDERMAL DRUG DELIVERY

Abstract

Protein-based nanocarriers are promising because they occur naturally and generally show less cytotoxicity than synthetic molecules. Nanocarriers have been explored for drug delivery, especially in chemotherapy. They can target small pores, lower pH levels, and higher temperatures of cancer cells, potentially reducing the toxicity of many chemotherapeutic drugs. The skin protects the internal organs of human body with regard to the entry and exit of a large number of auxiliary products water salts and other metabolites. However, these barriers pose a challenge into the penetration of the drug into the skin and its diffusion across the membrane. Liposome preparations of anticancer drugs, antihypertensive drugs, steroidal and non-steroidal analgesic drugs, local ~~anesthetic~~[anaesthetic](#) drugs, antimigraine and antiparkinsonian drugs, vaccines, and toxoids are administered through transdermal delivery.

Keywords: cancer cells, chemotherapeutic drugs, antiparkinsonian drugs, transdermal delivery

Introduction:

Nanocarriers are nanomaterials used to transport substances, such as drugs. These carriers are now being utilized in drug delivery, demonstrating their potential in chemotherapy. Common nanocarriers include polymers, liposomes, micelles, polymeric nanoparticles, dendrimers, lipid-based carriers, carbon-based materials, and gold nanoparticles. Examples of gold nanoparticles include nanocages and gold nano-shells. Various nanomaterials in nanocarriers enable delivery of both hydrophobic and hydrophilic drugs throughout the body. Given that the human body is mostly water, the efficient delivery of hydrophobic drugs is a significant therapeutic advantage of nanocarriers. Micelles can incorporate either hydrophobic or hydrophilic drugs based on the orientation of phospholipid molecules. Nanocarriers are nanotube arrays that can accommodate both types of drugs.

However, a potential issue with nanocarriers is the undesirable toxicity of the nanomaterials that are used. Inorganic nanomaterials can be toxic if they accumulate in certain organelles. Further research is ongoing to develop more efficient and safer nanocarriers. [Protein-based nanocarriers are promising because they occur naturally and generally show less cytotoxicity than synthetic molecules.](#)

Comment [VP1]: Explain how and why?

Nanocarriers have been explored for drug delivery, especially in chemotherapy. They can target small pores, lower pH levels, and higher temperatures of cancer cells, potentially reducing the toxicity of many chemotherapeutic drugs. As approximately 75% of antitumor drugs are hydrophobic and difficult to deliver into human cells, using micelles to maintain and mask the hydrophobic nature of these drugs offers new prospects for hydrophobic anticancer drugs¹.

Drug delivery through the skin

Drugs applied to the skin can cross the epidermis through either the skin's appendages or the stratum corneum. The appendages cover only approximately 0.1% of the surface, so they contribute little to steady-state drug penetration, but they may be important for ions, large molecules, polymers, and colloidal particles. The stratum corneum, only 10-15 layers^w thick, is the main barrier for permeation. It has a "brick and mortar" structure with keratin-filled cells (bricks) in a matrix (mortar) made of long-chain ceramides, fatty acids, triglycerides, cholesterol, cholesterol sulfate, and sterol/wax esters. Proteins and desmosomes in the lipid layers add rigidity and resistance to diffusion. Most molecules pass through the lipid domains; therefore, most skin penetration enhancement methods focus on this route².

Table No. 1: Ideal properties of drugs for transdermal drug delivery³

S.No	Parameter	Properties
1	Dose	Should be low (less than 20mg/day)
2	Half-life	10 or less hours
3	Molecular weight	<400 Dalton
4	Partition coefficient	Log P (octanol -water) between 1.0 and 4.0
5	Skin permeability coefficient	$>0.5 \times 10^{-3}$ cm/h
6	Liophilicity	$10 < K_o/w < 1000$
7	Oral bioavailability	Low
8	Therapeutic window	Low
9	Melting point	$<200^{\circ}\text{C}$
10	pH	Between 5.0 to 9.0

Barriers of the delivery of the drug through skin

The skin protects the internal organs of human body with regard to the entry and exit of a large number of auxiliary products water salts and other metabolites. However, these barriers pose a challenge into the penetration of the drug into the skin and its diffusion across the

Comment [VP2]: Repeated sentence.....already mentioned In Introduction < Line 1

Comment [VP3]: Specify the range of temperature

Comment [VP4]: Give reason why?

Comment [VP5]: Which layer.....mentioned the thickness of skin in micrometer/nanometer/picometer.

10-15 layers of what?

Comment [VP6]: First explain what is lipid domain, why it is important in drug delivery and what kind of synergism molecules have with this?

Comment [VP7]: Why melting point of drug is important for transdermal drug delivery system?

What is the significance of melting point in TDDS?

Comment [VP8]: Title could be reframe as

"Challenges in Drug Delivery Through the Skin"

membrane. As the first layer of defense, the stratum corneum, topmost layer of the skin exposed to the environment, is usually considered a significant hurdle for a broad set of drugs. Highly lipophilic and small molecule drugs are easier to permeate but restricts other classes of compounds. Stratum corneum is mostly lined by corneocytes (to a large extent the keratin), and are responsible for the permeation of water during the standard homeostatic mechanisms⁴.

Interestingly, both dermal and transdermal drug delivery involve similar steps. The drug must first deposit on the stratum corneum and then diffuse into the viable epidermis. In transdermal delivery, the drug also needs to pass through the dermis to reach the systemic circulation⁵.

In dermal delivery, bypassing the stratum corneum to concentrate the drug in the epidermis sounds appealing, but it is rarely effective. The epidermis actively metabolizes drugs, and the drug may permeate into deeper layers, reducing its concentration within the skin. The drug might also diffuse into hair follicles and sweat glands, where it is further metabolized, decreasing transdermal permeation.




Figure 1: Routes of drug delivery through skin (1) through the appendageal route, (2) through the intracellular route, or (3) through the intercellular route⁴.

Several successful strategies have been developed to enhance drug permeation within the skin, including physical and chemical methods, their combination, and micro- and nano-technological approaches. Recently, polymeric, lipidic, and surfactant-based nano-drug delivery platforms have gained attention. These nano-systems can provide the needed lipophilicity, occlusion, irritation masking, and stabilization during diffusion, depending on the drug⁶. Most dermal and transdermal drug formulations use lipid-based systems due to the lipidic nature of skin cells and tissues. However, many lipid-based systems require polymeric

bases for application, enhancing permeation and delivery. There are various types of vesicular and non-vesicular transdermal drug delivery systems.

RATIONALE FOR TRANSDERMAL DRUG DELIVERY

The skin is an excellent barrier to molecular transport, so the rationale for this delivery strategy must be clear. There are times when oral drug intake isn't possible, and alternatives are needed. While intravenous (IV) drug administration avoids issues like gastrointestinal and liver metabolism, it is invasive and uncomfortable, especially for long-term use. This has led to the exploration of other delivery routes.

Advantages:

- Allows easy and pain-free self-administration for patients.
- Reduces dosing frequency and prevents the fluctuations in drug levels.
- Drugs with a short half-life can be delivered effectively.
- It improves patient compliance, especially for long-term treatments like chronic pain and smoking cessation⁷.
- It avoids the liver's first-pass metabolism and the GI tract for drugs with poor bioavailability.
- Transdermal systems are cheaper and can deliver drugs for 1 to 7 days.
- New programmable systems allow for multiple dosing and variable drug delivery, adding benefits to traditional patches.
- The transdermal route allows the use of potent drugs with minimal risk of system toxicity.
- In case of toxicity, the patient can simply remove the transdermal patch.

Disadvantages:

- Drugs needing high blood levels cannot be given this way.
- The adhesive may not stick well to all skin types.
- Drugs or formulations can cause skin irritation or allergies.
- Patches can be uncomfortable to wear.
- Manufacturing requires special equipment, expensive than traditional methods.
- There is a delay before the drug enters the bloodstream, so it's not good for drugs needing quick action⁸.

Elastic liposome:

Elastic liposomes (EL) are some of the most flexible deformable amphiphilic vesicular systems containing biocompatible lipids and surfactants for the diffusion of many

Comment [VP9]: Specify the types of molecules?

Comment [VP10]: Quote the reference?

Comment [VP11]: Justify with reference

Comment [VP12]: These two examples are very generic in today's time. Give some more examples following patents, press release and recent management of different disease and indication by TDDS.

Comment [VP13]: Very generic advantage

Comment [VP14]: Reference?

Comment [VP15]: Not clear what the author want to say?

Reframe with reference

Comment [VP16]: Reference?

For a potent drug the maximum oral dose range between 0.3mg to 20mg daily for different drugs. Then how would you administer such a low dose using TDDS which would be the more bioavailable then oral route? Justify your answer.

Comment [VP17]: What kind of toxicity occurred because of TDDS?

Comment [VP18]: What does it means.....please elaborate with reference

Comment [VP19]: Please explain all skin types

Comment [VP20]: Skin irritation studies always conducted during development.

With the help of some references explain when skin irritation was occurred due to the transdermal patch/ formulation.

difficult molecules for curative, biochemical, and cosmetic applications. It has been used in several pharmaceutical technologies through topical, transdermal, nasal, and oral delivery systems for better and enhanced drug delivery. ELs are similar to conventional liposomes but contain an edge activator in the lipid bilayer structure to make it flexible. ELs include deformable liposomes, ultra-deformable liposomes, flexible liposomes, ultra-flexible liposomes, and ~~transfersomes~~ transfersomes. They are composed of phospholipids and surfactants such as edge activators (EA), and they have an internal water core that is enclosed within a lipid layer and can contain hydrophilic materials in the aqueous chamber and lipophilic materials in the lipid bilayer. However, other materials such as charged lipids, including cationic and anionic lipids; polyethylene glycol, which is also known as PEG or PEGylation; ethanol; cyclodextrin complexes; and gels. Composition causes changes in physicochemical characteristics and, hence, skin permeability behavior, and therefore the efficacy of a chemical structure. For instance, it should be mentioned that it is a noticeable fact that in vivo skin permeation performance reflects quite different results compared to *in-vitro* data and cannot be explained by *in-vitro* results⁹.

Liposome preparations of anticancer drugs, antihypertensive drugs, steroidal and non-steroidal analgesic drugs, local anesthetic drugs, antimigraine and antiparkinsonian drugs, vaccines, and toxoids are administered through transdermal delivery (figure 2).




Figure 2: Structure of Liposomes

Niosomes:

Niosomes are vesicular nanocarriers. They possess lamellar (bilayer) structures composed of amphiphilic molecules with a surrounding aqueous phase (Fig. 3). Surfactants consist of both water-insoluble tails and water-soluble heads, which have self-assembling characteristics and form structures such as spheres or plate-like structures called micelles or lamellar bilayers. Some of the areas in surfactants that can be used in delivering drugs are

Comment [VP21]: Give me some examples of easy molecules.

Don't use terms like difficult molecules

Comment [VP22]: Here are you defining the elastic liposomes or the transfersomes? Make it clear.

Formatted: Font: Italic

Formatted: Font: Italic

Comment [VP23]: Is there any commercial product available as Liposome for different indication mentioned here.

sorbitan esters and analogs depending on sugar-based, polyoxyethylene, or polyglycerol or crown ether based nonionic surfactants are widely used as they avoid skin irritation followed by anionic and cationic surfactants¹⁰.

The structures of ~~niosomes~~~~noisome~~, which are considered vesicular systems, differ in their ability to entrap both hydrophilic and lipophilic materials. Hydrophilic drugs are either encapsulated in the inner aqueous core or adsorbed on bilayer surfaces, whereas lipophilic agents are incorporated into the lipid core of the bilayers through lipid solubility.

The preparation of vesicular assemblies depends on the supply of some form of energy; all examined techniques involve the hydration of one or more surfactants above the gel-to-liquid phase transition temperature of the system to form a colloidal dispersion. Owing to their possible capacity to encapsulate numerous chemotherapeutics, these vesicles are used as drug delivery systems to facilitate drug targeting, controlled release and permeation¹¹. ~~Niosomes~~ can act as therapeutic reservoirs for delivering a drug in an effective manner, which in turn increases the bioavailability of the drug, to obtain the therapeutic spectrum of the drug for an even longer period of time. The ~~niosomes~~~~noisome~~ can be altered in terms of components and composition, concentration of different additives, and the surface charge of components of the vesicles or membrane additive substances. Furthermore, it has been ascertained that drug ionization affects the physicochemical characteristics of vesicles as well as their percutaneous permeation profiles, and niosomes have been investigated in depth as potential carriers for sustained and targeted drug delivery, since they are easily derivatized to enhance vesicle versatility to improve the affinity for the target site¹².

Comment [VP24]: Reference?




Figure 3: Structure of Niosomes

Ethosomes:

Ethosomes are mucosal-targeted noninvasive preparations that allow drugs to ~~penetrate~~~~intopenetrate~~ the dermal or systemic circulation. They are soft, easily deformable, and highly compatible with the deposition of active agents. They mainly consist of phospholipids:

phosphatidylserine, phosphatidic acid, and high concentrations of ethanol (45%w/w) and water (Figure 4). Ethosomes are characterized by a high concentration of ethanol, which is known for its ability to disturb the organization of the skin lipid bilayer; therefore, penetration of such vesicles through the stratum corneum is possible. Second, because ethanol content has been loaded high in lipid membranes, but compared with common vesicles, the lipid membrane is looser, yet carries equal stability, which enhances malleability and improves the drug distribution performance in the lipids of the stratum corneum. The molecular weights of the drugs incorporated into classical ethosomes fell within the range of 130. Ethosomes are adaptable for different uses in the pharmaceutical/biotechnology, veterinary, cosmeceutical, and functional food/nutraceutical fields. These “soft vesicles” can be considered new vesicular carriers with improved delivery to and through the skin. Ethosome vesicles can be made to range between tens of nanometers to particles of the micron range¹³.

Comment [VP25]: Rephrase this line

Comment [VP26]: Of what?

Comment [VP27]: Mention.....what is the entrapment efficiency of Ethosomes in general?




Figure 4: Structure of Ethosomes

Transfersomes

Transfersomes, also called ultra-flexible liposomes or elastic vesicles, contain bilayers as their core and edge activators. Transfersomes are more sensitive to stress than are conventional liposomes. Therefore, the positive isotropic pressure and mixture of phospholipids and edge activators in an optimal ratio provide transfersomes with excellent deformability. These vesicles are relatively elastic in nature, primarily because of the action of the edge activator, which makes the lipoidal bilayer of vesicles flexible and deformable. Edge activators are single-chain surfactants with high radii of curvature. Spans, tweens, dipotassium glycyrrhizinate, sodium cholate, and sodium deoxycholate were the major edge activators (figure 5). When administered through the transdermal route, transfersomes expand and enter deeper layers under the effect of a water gradient across the skin's horny layer, the stratum corneum, and settle in the subcutaneous tissue. Accumulated transfersomes help shield the built-in drug from cellular expulsion from the cutaneous

circulation and the release of drugs into the systemic circulation, leading to increased circulation time and drug bioavailability¹⁴.




Figure 5: Structure of Transfersomes¹⁵

Invasome

Invasomes are also a modern vesicular system and are of vital importance for enhancing the penetration of active molecules of drugs through the skin as compared to other vesicles. These vesicles are formed from phospholipids, ethanol, terpenes, or a blend of two or more terpenes. These components have proven to be good transdermal penetrators with the required penetration characteristics. Invasomes are soft liposomal vesicles containing insignificant amounts of ethanol and terpene or terpene combinations and are potential transporters that exhibit enhanced skin permeability. These lipidic carriers are consisted of phospholipids i.e. phosphatidylcholine, phosphatidylserine, soya phospholipid, egg lecithin, phosphatidylinositol, phosphatidic acid and phosphatidylglycerol), low concentration of ethanol (3%–3.3% v/v), terpenes or a mixture of terpenes (i.e. citral, cineole, limonene, eugenol; 1–5% v/v) and water (Figure 5). Terpenes have the general formula $(C_5H_8)_n$, which enhances the rates of skin permeation of both hydrophilic and hydrophobic drugs. Terpenes are constituents of plant-derived essential oils and are used extensively as carriers or envisaged penetration enhancers. However, terpenes have other advantages, such as low skin irritation at lower concentrations. Moreover, according to FDA classes, terpenes are among the products generally considered safe to use¹⁶.

Comment [VP28]: Mention.....what is their entrapment efficiency, how much flexible they are, how much they can squeeze themselves???

Comment [VP29]: Reference?

Comment [VP30]: Mention.....what is their entrapment efficiency, how much flexible they are, how much they can squeeze themselves???




Figure 5: Structure of Invasome

Glycosomes:

Glycosomes are bilayer structures used for dermal and transdermal drug delivery. Glycerol is a viscous liquid and an alcohol. It has three hydroxyl groups that make it naturally soluble in water, making it a polar molecule. Glycerol is a triglyceride present in animal fat and vegetable oils. This can be made directly ~~during the course of~~during soap production, or it can be a co-product derived from biodiesel production. It is used in the formulation of drugs as lubricants, humectant agents, edge promoters, and emulsifying agent. These vesicles are quite dissimilar to the common liposomes ~~with regard to~~regarding bilayer fluidity prepared by the incorporation of phospholipids and different concentrations of glycerol (10-30 % v/v). These vesicles effectively transport active ingredients across the skin tissue. Liposomes are less stable and less fluid than glycosomes and are therefore commonly used in topical drug delivery systems. This is a relatively noninvasive and non-toxic procedure for drug administration. Because glycerol increases the deformability index of liposomal bilayers, there is better penetration into the skin. Liposomes, which are formed by phospholipids and ~~cholesterol,~~ glycosomes~~cholesterol,~~ glycosomes are new vesicular systems. If phospholipids are dispersed in water, they rapidly self-organize into bilayer vesicle structures¹⁷.

Cubosomes:

Cubosomes are a type of nanoparticle, but unlike solid particles, they are self-assembled liquid crystalline particles formed from certain surfactants mixed with water in the right ratio (**Figure 6**). Their unique microstructure offers properties of significant practical interest. The discovery of cubosomes is intriguing and involves aspects of food science, differential geometry, biological membranes, and digestive processes. One of the most commonly used surfactants to create cubosomes is the monoglyceride glycerol monoolein. The bicontinuous cubic liquid crystalline phase, known for being optically clear and highly viscous, has a

Comment [VP31]: Mention.....what is their entrapment efficiency, how much flexible they are, how much they can squeeze themselves???

Figure??

distinctive nanometer-scale structure. The term "bicontinuous" describes the separation of two continuous, non-intersecting aqueous regions by a lipid bilayer twisted into a space-filling form. Cubosomes are typically formed by hydrating a surfactant or polar lipid that forms a cubic phase and then dispersing the solid-like phase into smaller particles. There is significant interest in cubic phases due to their biologically compatible microstructure and ability to control the release of solubilized active ingredients, such as drugs and proteins¹⁸




Figure 6: Structure of Cubosomes¹⁸

References:

1. Namdari, M., Eatemadi, A., Soleimaninejad, M., &Hammed, A. T. (2017). A brief review on the application of nanoparticle enclosed herbal medicine for the treatment of infective endocarditis. *Biomedicine & Pharmacotherapy*, 87, 321-331.
2. Krishna Reddy Y, Maheswara Reddy D, Ashok Kumar B. Transdermal drug delivery system. *Indian journal of Pharmacy and Biotechnology*. 2014;2(2):1094-1103.
3. Patel BA. Permeationenhancement and advanced strategies: a comprehensive review of improved topical drug delivery. *International Research Journal of Modernization in Engineering Technology and Science*. 2024;6(05):6691-702.
4. Palmer BC, DeLouise LA. Nanoparticle-enabled transdermal drug delivery systems for enhanced dose control and tissue targeting. *Molecules*. 2016 Dec 15;21(12):1719.
5. Matharoo N, Mohd H, Michniak-Kohn B. Transfersomes as a transdermal drug delivery system: Dermal kinetics and recent developments. *Wiley Interdisciplinary Reviews: Nanomedicine and Nanobiotechnology*. 2024 Jan;16(1):e1918.

Comment [VP32]: Mention.....what is their entrapment efficiency, how much flexible they are, how much they can squeeze themselves???

Formatted: Font: 12 pt, Danish (Denmark)

6. Rabiei M, Kashanian S, Samavati SS, Jamasb S, McInnes SJP. Nanomaterial and advanced technologies in transdermal drug delivery. *J Drug Target*. 2020;28:356–367.
7. Chunyunzhu, Yongtaizhang, Tong wu, Zehui he, Teng guo, Nianping feng. Optimizing glycosome formulations via an orthogonal experimental design to enhance transdermal triptolide delivery, *Acta Pharm*. 2022;72:135–146.
8. Deulkar DA, Kubde JA, Hatwar PR, Bakal RL. A review on transdermal drug delivery system. *GSC Advanced Research and Reviews*. 2024;18(2):347-61.
9. Sinico C, Fadda AM. Vesicular carriers for dermal drug delivery. *Expert Opin Drug Deliv*. 2009;6(8):813–825.
10. Moghassemi S, Hadjizadeh A. Nano-niosomes as nanoscale drug delivery systems: an illustrated review. *J Control Release*. 2014;185:22–36.
11. Gannu PK, Rajeshwarrao P. Nonionic surfactant vesicular systems for effective drug delivery an overview. *Acta Pharmacol Sin*. 2011;1:208–219.
12. Perche F, Torchilin VP. Recent trends in multifunctional liposomal nanocarriers for enhanced tumor targeting. *J Drug Deliv*. 2013;2013:705265.
13. Shinde P, Page A, Bhattacharya S. Ethosomes and their monotonous effects on skin cancer disruption. *Frontiers in nanotechnology*. 2023;12(5):1087413.
14. Simrah, Hafeez A, Usmani SA, Izhar MP. Transfersome, an ultra-deformable lipid-based drug nanocarrier: an updated review with therapeutic applications. *Naunyn-Schmiedeberg's Archives of Pharmacology*. 2024 Feb;397(2):639-73.
15. Rai S, Pandey V, Rai G. Transfersomes as versatile and flexible nano-vesicular carriers in skin cancer therapy: The state of the art. *Nano reviews & experiments*. 2017 Jan 1;8(1):1325708.
16. Singh D. Ultradeformable system: a carrier for transdermal drug delivery. *IJPSR*. 2013;4: 4098–4104.
17. Manca ML, Zarub M, Manconi M, Lai F, Valenti D, Sinico C, et al. Glycosomes: A new tool for effective dermal and transdermal drug delivery. *Int J Pharm*. 2013;455:66-74
18. Umar H, Wahab HA, Gazzali AM, Tahir H, Ahmad W. Cubosomes: Design, Development, and Tumor-Targeted Drug Delivery Applications. *Polymers*. 2022; 14(15):3118.
19. Chacko IA, Ghate VM, Dsouza L, Lewis SA. Lipid vesicles: A versatile drug delivery platform for dermal and transdermal applications. *Colloids and Surfaces B: Biointerfaces*. 2020 Nov 1;195:111262.
20. Rastogi V, Yadav P. Transdermal drug delivery system: An overview. *Asian Journal of Pharmaceutics (AJP)*. 2012;6(3).

Formatted: Font: (Default)
TimesNewRomanPS, 12 pt, Font color: Black,
Danish (Denmark)

21. Muzzalupo R, Tavano L. Niosomal drug delivery for transdermal targeting: recent advances. *Research and reports in transdermal drug delivery*. 2015 Jul 29:23-33.
22. Namdari M, Eatemadi A, Soleimanejad M, Hammed AT. A brief review on the application of nanoparticle enclosed herbal medicine for the treatment of infective endocarditis. *Biomedicine & Pharmacotherapy*. 2017 Mar 1;87:321-31.

UNDER PEER REVIEW