

# Theoretical study of Properties of Indoline-2,3-dione-3-oxime using Chemicalize Software

## Abstract:

Indoline-2,3-dione-3-oxime (IDOX) (Isatin -3-oxime) is employed as a anticonvulsant and is known for treating various disorders caused by release of amino acids responsible for anxiety and excitatory behavior. In view of the biological importance of IDOX Properties, pka, Isoelectric Point, logP, logD, Solubility, Geometry, HNMR were determined by using Chemicalize software

Abstract is too short which will not meet the journal requirement, abstract should not be less than 250 words

**Keywords:** Indoline-2,3-dione-3-oxime, Properties, pka, Isoelectric Point, logP, logD, Solubility, Geometry, HNMR

## Introduction

In disorders such as cerebral ischemia or cerebral infarction leading to bad effects like hemorrhagic stroke, cardiac arrest and hypoglycemia excitatory amino acids cause to the degeneration of neurons. Compounds like Indoline-2,3-dione-3-oxime (Isatin-3-oxime)<sup>1-4</sup> possess property of blocking excitatory amino acid receptors which have therapeutic use over the effects of above breakdown [29].

From the Literature studies it is evident anticonvulsant screening of Indoline-2,3-dione-3-oxime (IDOX) was carried out by applying Maximal Electroshock (MES) model and the results indicated that IDOX is active at lower dose of 100 mgkg<sup>-1</sup>. It is also found that IDOX is more potent when compared to standard drug sodium valproate [31].

In appropriate citation and format

With the application of chemicalise software the structure of Indoline-2,3-dione-3-oxime (IDOX) is drawn and results obtained were interpreted in this present paper. Basic properties, structural properties, names and identifiers of IDOX were given in this present paper. The details of pKa, isoelectric point, logP, logD, Solubility, Geometry and <sup>1</sup>H NMR spectrum of IDOX were also studied in this manuscript [30].




Fig.1 structure of Indoline-2,3-dione-3-oxime (IDOX)

## Hypothetical Study of Indoline-2,3-dione-3-oxime using Chemicalize Software

### Basic properties of Indoline-2,3-dione-3-oxime (IDOX)

Indoline-2,3-dione-3-oxime (IDOX) is drawn as the input of Chemicalize Software<sup>5,6</sup> and from the data obtained (Table.1) it is evident that the formula of Indoline-2,3-dione-3-oxime is  $C_8H_6N_2O_2$ . The data obtained also indicated that composition of Indoline-2,3-dione-3-oxime (IDOX) is C=59.26%, H=3.73%, N=17.28%, S (19.73%).

Mention about the software details and also highlights discuss about IDOX

Table 1. properties of Indoline-2,3-dione-3-oxime (IDOX) as determined by chemicalize software

Input	Indoline-2,3-dione-3-oxime
Use major tautomer	✓ (the input is the major tautomer)
Molar mass	162.148 g/mol
Exact mass	162.042927441 Da
Formula	$C_8H_6N_2O_2$
Composition	C (59.26%), H (3.73%), N (17.28%), O (19.73%)
Lipinski's rule of five	✓

### Structural Properties of Indoline-2,3-dione-3-oxime (IDOX)

Structural calculations of Indoline-2,3-dione-3-oxime (IDOX) as indicated by Chemicalise software (Table.2) enabled the evaluation of various topology-related characteristics, hydrogen bonding, and other physical properties. Table.4 shows the Data of structural properties of Indoline-2,3-dione-3-oxime (IDOX).

In Table 4. Mention information is not appropriate

Chemical formula of IDOX is  $C_8H_6N_2O_2$  and this in agreement with the atom count of 18 and this confirms the Chemical formula. The total heavy atom count of 12 which indicates the presence of 8 carbons, 2 nitrogens and 2 oxygens in IDOX (fig 1&2). There is no asymmetric atom in IDOX and has no rotatable bonds<sup>7-8</sup>. Chemicalize data of IDOX indicates the presence of two rings where one is aromatic ring and other one is hetero ring. H bond donor count is 2 and H bond acceptor count is 3 in IDOX. The topological polar surface area<sup>10,11</sup> of Indoline-2,3-dione-3-oxime is  $61.69 \text{ \AA}^2$ , polarizability is  $15.91 \text{ \AA}^3$  and its Molar refractivity is indicated to be  $44.3 \text{ cm}^3/\text{mol}$

**Table 2. Structural properties of Indoline-2,3-dione-3-oxime as determined by Chemicalize**

UNDER PEER REVIEW

## Structural properties

Atom count	18
Heavy atom count	12
Asymmetric atom count	0
Rotatable bond count	0
Ring count	2
Aromatic ring count	1
Hetero ring count	1
FSP3	0
Hydrogen bond donor count	2
Hydrogen bond acceptor count	3
Formal charge	0
Topological polar surface area	61.69 Å <sup>2</sup>
Polarizability	15.91 Å <sup>3</sup>
Molar refractivity	44.3 cm <sup>3</sup> /mol

### Names and Identifiers of Indoline-2,3-dione-3-oxime(IDOX)

IUPAC name of IDOX is 3-(hydroxyimino)-2,3-dihydro-1H-indol-2-one and its the commonly used trivial or traditional name is isatin-3-oxime. other details like SMILES, InChI and InChIKey of IDOX as given by the chemcalise software are given in Table 3.


Table 3. Names and identifiers of Indoline-2,3-dione-3-oxime as given by chemicalize software

Names and identifiers	
IUPAC name	3-(hydroxyimino)-2,3-dihydro-1H-indol-2-one
Traditional name	isatin-3-oxime
Common names	N/A
SMILES	<chem>ON=C1C(=O)NC2=CC=CC=C12</chem>
InChI	InChI=1S/C8H6N2O2/c11-8-7(10-12)5-3-1-2-4-6(5)9-8/h1-4,12H,(H,9,10,11)
InChIKey	LNMAXZZQNSPQSR-UHFFFAOYSA-N


### pKa of Indoline-2,3-dione-3-oxime

pKa values of IDOX are found to be 7.13 and 15.31 as indicated by Chemcalise data<sup>12-16</sup> (fig.2). pKa value of 7.13 corresponds to dissociation of H of =N-OH group of IDOX and this indicates neutral nature of this group. pKa value of 15.31 corresponds to dissociation of imino H atom of -NH-CO group. This attributes to the strong basic nature of Indoline-2,3-dione-3-oxime (IDOX).

**Fig.2 Distribution of pH for the different microspecies of Indoline-2,3-dione-3-oxime as given by chemicalize software**



Missing fig.No. and Name




Missing fig.No. and Name

Isoelectric Point of Indoline-2,3-dione-3-oxime

Distribution of charge with pH of Indoline-2,3-dione-3-oxime as derived using chemicalize software is given in figure 3. The data of distribution of charge with pH of IDOX indicates that its isoelectric point is 2.8. IDOX is electrically neutral with zero electric charge within the pH range of 1.7 to 4.6 thereby indicating the isoelectric point<sup>17,18</sup> of 2.8.

In fig3. PH of IDOX isoelectric point 2.8 is missing and not justified

**Fig.3** Distribution of charge with pH of Indoline-2,3-dione-3-oxime as given by chemicalize software




## log P and log D of Indoline-2,3-dione-3-oxime

logP value of Indoline-2,3-dione-3-oxime<sup>19-21</sup> is found to be 0.961.

Variation of the log D value of IDOX with pH is given in figure 4. log D value of IDOX is 0.96 within the pH range of 1.7 to 4.6 and further as the pH increases log D value decreases which is found to be 0.05 at pH 8 (Fig 4).

**Fig.4 Distribution of log D with pH of Indoline-2,3-dione-3-oxime as given by chemicalize software**




## Solubility (log S) of Indoline-2,3-dione-3-oxime(IDOX)

Distribution of log S with pH of IDOX as obtained by using Chemicalise software is given in Figure.5 and the data indicates that Indoline-2,3-dione-3-oxime<sup>22-25</sup> is of High solubility category with Intrinsic solubility of -1.589.

Elaborate the solubility and the given data and the graph is not appropriate and justified its varying

Fig.5 Distribution of log S with pH of Indoline-2,3-dione-3-oxime as given by chemicalize software



## Geometry of Indoline-2,3-dione-3-oxime(IDOX)



Various geometrical parameters like vanderwaals volume, vanderwaals surface area, solvent accessible surface area, topological polar surface area, minimum and maximum projection area, minimum and maximum projection radius of Indoline-2,3-dione-3-oxime(IDOX) were measured with the application of chemicalize software<sup>26</sup> (Table. 4).

**Table. 4 Data of of volume, surface area ,projection area and projection radius of Indoline-2,3-dione-3-oxime as given by chemicalize software**

Geometry	
Van der Waals volume	130.9 Å <sup>3</sup>
Van der Waals surface area	197.92 Å <sup>2</sup>
Solvent accessible surface area	294.58 Å <sup>2</sup>
Topological polar surface area	61.69 Å <sup>2</sup>
Minimum projection area	25.15 Å <sup>2</sup>
Maximum projection area	54.07 Å <sup>2</sup>
Minimum projection radius	3.9 Å
Maximum projection radius	5.15 Å

## H-NMR Spectrum of Indoline-2,3-dione-3-oxime(IDOX)

NMR Predictor of Chemicalise software is employed for recording NMR spectra of Indoline-2,3-dione-3-oxime(IDOX). The estimation of chemical shifts<sup>27,28</sup> of IDOX is in relation to the chemical shift of tetra methylsilane (TMS = 0ppm). H NMR spectrum of IDOX is given in the Table 5.



**Table 5. shifts caused by the protons of Indoline-2,3-dione-3-oxime(IDOX) and the intensity and quality for each shift and corresponding protons as given by chemicalize software**

The shifts caused by the protons of Indoline-2,3-dione-3-oxime(IDOX) is given in Table 5. The intensity and quality for each shift of IDOX is also obtained from this data.1H NMR

spectrum of IDOX shows peaks at  $\delta$  9.82ppm (1H,=N-OH) and  $\delta$  6.77 ,7.11 , 7.159( 4H,Ar-H). The shift observed at 8.43 ppm also corresponds to protons of aromatic ring (Table.5 ).

## Conclusions

The present article reflects the application of Chemicalize software in understanding various structural and physicochemical aspects of Indoline-2,3-dione-3-oxime(IDOX).Data of chemicalize software from pKa values and <sup>1</sup>HNMR spectral data indicated that IDOX is of strong basic nature. Variation of properties like log D, log S, solubility in mg/L with pH of Indoline-2,3-dione-3-oxime(IDOX) is studied by Graphical interpretation by employing Chemicalize software. Apart from this the data of properties like vanderwaalsarea,vanderwaals volume, projection area projection radius of IDOX is also obtained by applying chemicalise software.

**Reference and citation is not up to the mark and not in proper format**

## References

1. Parvaneh Pakravan et al ,Biochemical and pharmacological characterization of isatin and its derivatives: from structure to activity Pharmacological Reports 2013,65,313-335
2. Smitha. S, Pandeya .S. N., *Sci. Pharm.* 2008, 76, 621.
3. Pandeya. S. N, Kumar. R, *Asia J. Res. Chem.*, 2010, 3(3): 646.
4. Pandeya S. N, *IJPI's Journal of Drug Chemistry*2012 , 2:8,1-7
5. G. Audi, A.H.Wapstra, The 1995 update to the atomic mass evaluation, Nuclear Physics A, Vol. 595, Issue 4, pp. 409- 480, 1995.
6. Trepalin, S.V., Bessonov, Y.E., Fel'dman, B.S. et al. The Structural Chemical Database of the All-Russian Institute for Scientific and Technical Information, Russian Academy of Sciences. An Autonomous System for Structural Searches. *Autom. Doc. Math. Linguist.* 52, 297–305 (2018). <https://doi.org/10.3103/S0005105518060055>
7. Molecules from the Minkowski space: an approach to building 3D molecular structures,December 2003,Journal of Molecular Structure THEOCHEM s 666–667(6C):51–59 DOI:10.1016/j.theochem.2003.08.013
8. Bender, B.J., Gahbauer, S., Lutten, A. et al. A practical guide to large-scale docking. *Nat Protoc* 16, 4799–4832 (2021). <https://doi.org/10.1038/s41596-021-00597-z>
9. ChemAxon poster: Advanced automatic generation of 3D molecular structures
10. Ferrara P, Apostolakis J, Caflisch A. Evaluation of a fast implicit solvent model for molecular dynamics simulations. *Proteins.* 2002 Jan 1;46(1):24-33. doi: 10.1002/prot.10001. PMID: 11746700.
11. Norma Flores-Holguín<sup>1</sup>, Juan Frau<sup>2</sup>, Daniel Glossman-Mitnik,Computational Chemistry and Molecular Modeling Techniques for the Study of Micropeptin EI-964: Insights into Its Chemical Reactivity and Potential Pharmaceutical Properties, *Computational Molecular Bioscience*, Volume 13, Number 3, September 2023 ,DOI: 10.4236/cmb.2023.133003
12. Theoretical Determination of the pKa Values of Betalamic Acid Related to the Free Radical Scavenger Capacity: Comparison Between Empirical and Quantum Chemical Methods,MarcoTutone, Antonino Lauria, Anna Maria Almerico, July 2015,Interdisciplinary

13. Calculating pKa values of small and large molecules, J.Szegezdi and F.Csizmadia , American Chemical Society Spring meeting ,March 25-29th,2007
14. Epik: A software program for pKa prediction and protonation state generation for drug-like molecules,,January 2008,,Journal of Computer-Aided Molecular Design 21(12):681-91  
DOI:10.1007/s10822-007-9133-z
15. Epik version 2.0 (2009) Schrodinger, LLC, New York
16. Determination of acid dissociation constants (pKa) of cephalosporin antibiotics: Computational and experimental approaches,Chemosphere, Alyson R. Ribeiro, Torsten C. Schmidt,Volume 169, 2017,Pages 524-533,  
<https://doi.org/10.1016/j.chemosphere.2016.11.097>.
17. Calculation of the isoelectric points of native proteins with spreading of pKavalues,GHenriksson 1, A K Englund, G Johansson, P Lundahl, Electrophoresis , 1995  
Aug;16(8):1377-80,doi: 10.1002/elps.11501601227
18. plChemiSt – Free Tool for the Calculation of Isoelectric Points of Modified Peptides, Andrey I Frolov , Sunay V Chankeshwara, ZeyedAbdulkarim , Journal of Chemical information and Modelling, 2023 Jan 9;63(1):187-196. doi: 10.1021/acs.jcim.2c01261. Epub 2022 Dec 27.
19. Bouchard G., Carrupt P.A., Testa B., Gobry V., Girault H.H. Pharmaceutical Research 18 (2001) 702-708. <https://doi.org/10.1023/A>
20. Bouchard G, Pagliara A, Carrupt PA, et al. Theoretical and experimental exploration of the lipophilicity of zwitterionic drugs in the 1,2-dichloroethane/water system. Pharmaceutical Research. 2002 Aug;19(8):1150-1159. DOI: 10.1023/a:1019846125723. PMID: 12240941.
21. Işık, M., Bergazin, T.D., Fox, T. et al. Assessing the accuracy of octanol–water partition coefficient predictions in the SAMPL6 Part II log P Challenge. J Comput Aided Mol Des 34, 335-370 (2020). <https://doi.org/10.1007/s10822-020-00295-0>
22. Völgyi, G.; Baka, E. et al. Study of pH-dependent solubility of organic basis. Revisit of the Henderson-Hasselbalch relationship, AnalyticaChimicaActa, 2010 , 673, 40-46
23. Avdeef, A. et al. Equilibrium solubility measurement of ionizable drugs - consensus recommendations for improving data quality, ADMET & DMPK 4(2), 2016 , 117-178
24. Shoghi, E.; Fuguet, E.; Bosch, E.; Rafols, C. Solubility-pH profiles of some acidic, basic and amphoteric drugs, European Journal of Pharmaceutical Sciences 2013 , 48, 291-300
25. Alex Avdeef ,Suggested Improvements for Measurement of Equilibrium Solubility-pH of IonizableDrugs,July 2015,ADMET & DMPK 3(2):84-109,DOI:10.5599/admet.3.2.193
26. Chemaxon article: Molecules from the Minkowski space: an approach to building 3D molecular structures
27. . J Org Chem. 1997 Oct 17;62(21):7512-7515. doi: 10.1021/jo971176v. Authors. Hugo E. Gottlieb , Vadim Kotlyar, Abraham Nudelman
28. NMR Chemical Shifts of Common Laboratory Solvents as Trace Impurities Hugo E. Gottlieb, Vadim Kotlyar, and Abraham Nudelman J. Org. Chem., 1997, 62 (21), 7512-7515 • DOI: 10.1021/jo971176v Downloaded from http://pubs.acs.org on January 23, 2009  
IJRAR24A1683 207 International

29. KATANGURU LAXMI. Characterization of Keto and Enol Forms of Indoline-2,3-dione-3-oxime using Avogadro software Tool (A Theoretical Study). *Orient. J. Chem.*, Vol. 34(3), 1270-1277 (2018).
30. K. Laxmi. Calculation of Properties of Phenobarbital an Antiepileptic Drug Using Chemcalise Software. *International Journal of Scientific and Research Publications*, Volume 9, Issue 11, November 2019.
31. K. Laxmi. A study on structural aspects of indoline-2, 3-dione-3-oxime: Experimental and theoretical approach. *International Journal of Computational and Theoretical Chemistry*. 2013; 1(2): 11-17.

UNDER PEER REVIEW